

濱口先生
Preparation of Immunotolerant Space Under the Skin and Transplantation of Islets in the Space. <i>Tissue. Eng. Part. A.</i> 25(3-4): 183-192, 2019. こちらに邦題の入力をお願いいたします。
Prevalence and factors associated with sarcopenia in patients with rheumatoid arthritis. <i>Mod. Rheumatol.</i> 29(4): 589-595, 2019.
The Effects of Well-Rounded Exercise Program on Systemic Biomarkers Related to Cartilage Metabolism. <i>Cartilage</i> 10(4): 451-458, 2019.
牛込先生
Effects of dietary salt restriction on home blood pressure in diabetic patients with excessive salt intake: a pilot study. <i>J. Clin. Biochem. Nutr.</i> 65(3): 252-257, 2019.
Evaluation of the efficacy of simplified nutritional instructions from physicians on dietary salt restriction for patients with type 2 diabetes mellitus consuming excessive salt: protocol for a randomized controlled trial. <i>Trials</i> 23 20(1): 761, 2019.
橋本先生
Sarcopenia is associated with tongue pressure in older patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort study. <i>Geriatr. Gerontol. Int.</i> 19(2): 153-158, 2019.
Frequent Usage of Convenience Stores is Associated with Low Diet Quality. <i>Nutrients</i> 11(6): 1212, 2019.
岡村先生
Protein intake is not associated with progression of diabetic kidney disease in patients without macroalbuminuria. <i>Diabetes. Metab. Res. Rev.</i> 35(5): e3150, 2019.
The sodium-glucose cotransporter 2 inhibitor luseogliflozin can suppress muscle atrophy in Db/Db mice by suppressing the expression of foxo1. <i>J. Clin. Biochem. Nutr.</i> 65: 23-28, 2019.
High brain natriuretic peptide is associated with sarcopenia in patients with type 2 diabetes: a cross-sectional study of KAMOGAWA-DM cohort study. <i>Endocr. J.</i> 66: 369-377, 2019.
Triglyceride-glucose index is a predictor of incident chronic kidney disease: a population-based longitudinal study. <i>Clin. Exp. Nephrol.</i> 23: 948-955, 2019.
Maximum morning home systolic blood pressure is an indicator of the development of diabetic nephropathy: The KAMOGAWA-HBP study. <i>J. Diabetes. Investig.</i> 10: 1543-1549, 2019.
Short sleep duration is a risk of incident nonalcoholic fatty liver disease: a population-based longitudinal study. <i>J. Gastrointestin. Liver. Dis.</i> 28(1): 73-81, 2019.
Ectopic fat obesity presents the greatest risk for incident type 2 diabetes: a population-based longitudinal study. <i>Int. J. Obes.</i> 43: 139-148, 2019.
Shortage of energy intake rather than protein intake is associated with sarcopenia in elderly patients with type 2 diabetes: A cross-sectional study of the KAMOGAWA-DM cohort. <i>J. Diabetes.</i> 11(6): 477-483, 2019.
北江先生

The Triglyceride and Glucose Index Is a Predictor of Incident Nonalcoholic Fatty Liver Disease: A Population-Based Cohort Study. *Can. J. Gastroenterol. Hepatol.* 2019.

木村先生

Japanese radio calisthenics prevents the reduction of skeletal muscle mass volume in people with type 2 diabetes. *BMJ Open Diabetes Research and Care*: 10. 1136/bmjdr-2019-001027. 2019.